

## Publikationen 2018

(zuletzt aktualisiert: 01.02.2019)



Adelhardt T, Hessemeyer S, Heckel M, **Herbst FA, Stiel S**, Ostgathe C, Schöffski O. Kosten von multiresistenten Erregern in der stationären Palliativversorgung. Gesundh ökon Qual manag. 2018; DOI: 10.1055/a-0651-6448

**Afshar K**, Bleidorn Jutta, Büttner K, Schneider N, Steffens S. Kartierung kompetenzbasierter Lernziele im Curriculum Allgemeinmedizin an der Medizinischen Hochschule Hannover. Z Allg Med. 2018; 10: DOI: 10.3238/zfa.2018.0405-0411

**Afshar K**, Feichtner A, Boyd K, Murray S, **Jünger S, Wiese B, Schneider N, Müller-Mundt G**. Systematic development and adjustment of the German version of the Supportive and Palliative Care Indicators Tool (SPECT-DE). BMC Palliat Care. 2018; 17: 27

**Afshar K**, Fleischmann N, Schmiemann G, **Bleidorn J**, Hummers-Pradier E, Friede T et al. Reducing antibiotic use for uncomplicated urinary tract infection in general practice by treatment with uva-ursi (REGATTA) – a double-blind, randomized, controlled comparative effectiveness trial. BMC Complement Altern Med. 2018; 18: 23

Bock JO, Hajek A, Lühmann D, Ernst A, **Mamone S, Wiese B** et al. Unterschiede in Kognition, Depressivität und gesundheitsbezogener Lebensqualität zwischen Versicherten der GKV und PKV im hohen Alter. Psychiatr Prax 2018; 45: 148-153

Brettschneider C, Hajek A, Stein J, Luck T, **Mamone S, Wiese B** et al. Informelle Pflege bei Demenz nach Leistungsarten. Nervenarzt 2018; 89: 509-515

Dorow M, Stein J, Pabst A, Weyerer S, Werle J, Maier W, Miebach L, Scherer M, Stark A, **Wiese B, Moor L** et al. Categorical and dimensional perspectives on depression in elderly primary care patients - Results of the AgeMooDe study. Int J Methods Psychiatr Res 2018; 27

**Ewertowski H, Tetzlaff F, Stiel S, Schneider N, Jünger S**. Primary palliative Care in General Practice – study protocol of a three-stage mixed-methods organizational health services research study. BMC Palliat Care. 2018; 17(1): 21

Exner A, Kleinstäuber M, Maier W, Fuchs A, Petersen JJ, Schäfer I, Gensichen J, Riedel-Heller SG, Weyerer S, Bickel H, König H-H, **Wiese B** et al. Cross-cultural validation of the German version of the Four-Dimensional Symptom Questionnaire (4DSQ) in multimorbid elderly people. Qual Life Res. 2018; DOI: 10.1007/s11136-018-1924-2

Fischer K, Melo van Lent D, Wolfsgruber S, Weinhold L, Kleineidam L, Bickel H, Scherer M, Eisele M, van den Bussche H, **Wiese B** et al. Prospective Associations between Single Foods, Alzheimer's Dementia and Memory Decline in the Elderly. *Nutrients*. 2018;10(7): 852

Gerhard C, Habig H, Hagen O, Heusinger von Waldegg G, Knopf B, **Müller-Mundt G** et al. DGP-AG Nichttumorpatienten: Frühe palliative Versorgung von Menschen mit Nichttumorerkrankungen. *Z Palliativmed*. 2018; 19: 226-232

Hager K, Temps T, **Krause O**. Antidementiva, Antidepressiva und Neuroleptika bei alten Patienten absetzen. *Med Klin Intensivmed Notfmed*. 2018; DOI: 10.1007/s00063-018-0451-9

Hajek A, Brettschneider C, van den Bussche H, Kaduszkiewicz H, **Oey A, Wiese B** et al. Longitudinal Analysis of Outpatient Physician Visits in the Oldest Old: Results of the AgeQualiDe Prospective Cohort Study. *J Nutr Health Aging*. 2018; 22(6): 689-694

Hajek A, Brettschneider C, van den Bussche H, Lühmann D, **Oey A, Wiese B** et al. Impact of falls on depressive symptoms among the oldest old: Results from the AgeQualiDe study. *Int J Geriatr Psychiatry*. 2018; 33(10): 1383-1388

Heckel M, **Stiel S, Herbst FA**, Tiedtke JM, Sturm A et al. Multidrug-resistant bacterial microorganisms (MDRO) in end-of-life care: development of recommendations for hospitalized patients using a mixed-methods approach. *Support Care Cancer*. 2018; 26(9): 3021-3027

**Herbst FA**, Heckel M, Tiedtke JM, Adelhardt T, Sturm A, **Stiel S** et al. Perspectives on multidrug-resistant organisms at the end of life: A focus group study of staff members and institutional stakeholders. *Z Gerontol Geriatr*. 2018; DOI: 10.1007/s00391-018-1378-8

Heser K, Luck T, Röhr S, **Wiese B**, Kaduszkiewicz H, **Oey A** et al. Potentially inappropriate medication: Association between the use of antidepressant drugs and the subsequent risk for dementia. *J Affect Disord*. 2018; 226: 28-35

Heser K, Stein J, Luppa M, **Wiese B, Mamone S**, Weyerer S et al. Late-life depressive symptoms are associated with functional impairment cross-sectionally and over time: Results of the AgeMooDe study. *J Gerontol B Psychol Sci Soc Sci*. 2018; DOI: 10.1093/geronb/gby083

**Junius-Walker U**, Onder G, Soleymani D, **Wiese B**, Albaina O, Bernabei R et al. The essence of frailty: A systematic review and qualitative synthesis on frailty concepts and definitions. *European Journal of Internal Medicine*. 2018; DOI: <https://doi.org/10.1016/j.ejim.2018.04.023>.

Klein C, Wittmann C, Wendt KN, Ostgathe C, **Stiel S**. Palliative Sedierung. *Anaesthesist* 2018; DOI: 10.1007/s00101-018-0451-4

Kleineidam L, Thoma MV, Maercker A, Bickel H, Mösch E, Hajek A, König HH, Eisele M, Mallon T, Luck T, Röhr S, Weyerer S, Werle J, Pentzek M, Fuchs A, **Wiese B, Mamone S** et al. What Is Successful Aging? A Psychometric Validation Study of Different Construct Definitions. *Gerontologist*. 2018; DOI: 10.1093/geront/gny083

Knudsen L, Nawrotzki R, Schmiedl A, Mühlfeld C, **Kruschinski C**, Ochs M. Hands-on or no hands-on training in ultrasound imaging: A randomized trial to evaluate learning outcomes and speed of recall of topographic anatomy. *Anat Sci Educ.* 2018; 11(6): 575-591

**Krause O**, Hensen J. Säure-Basen-Haushalt. In: Blum HE, Müller-Wieland D (Eds.) *Klinische Pathophysiologie* (begründet von Walter Siegenthaler). Stuttgart: Thieme-Verlag; 2018. S. 288-300

**Krause O, Schleef T, Raker M, Schneider N, Bleidorn J.** Treatment of ambulant patients by a general practitioner within a university hospital's emergency department – follow-up study of patients' behaviour shortly afterwards. *GMS Ger Med Sci.* 2018; 16: Doc02; DOI: 10.3205/000264

Laidig F, May M, Brinkmann J, **Schneider N**, Stichtenoth DO. Evaluation of potential prescribing errors in patients with polypharmacy: a method to improve medication safety in ambulatory care. *Drugs Ther Perspect.* 2018; 34: 322-334

Mallon T, Ernst A, Brettschneider C, König H-H, Luck T, Röhr S, Weyerer S, Werle J, Mösch E, Weeg D, Fuchs A, Pentzek M, Kleineidam L, Heser K, Riedel-Heller S, Maier W, **Wiese B** et al. Prevalence of pain and its associated factors among the oldest-olds in different care settings - results of the AgeQualiDe study. *BMC Fam Pract.* 2018; 19: 85

McIntosh J, Alonso A, MacLure K, Stewart D, Kempen T, Mair A, Castel-Branco M, Codina C, Fernandez-Llimos F, Fleming G, Gennimata D, Gillespie U, Harrison C, Illario M, **Junius-Walker U** et al. A case study of polypharmacy management in nine European countries: Implications for change management and implementation. *PLoS One.* 2018; 13: e0195232

Poggenburg S, Schirgi J, **Afshar K**, Müller B, Bouchi-Häfner J, Siebenhofer A. Rückblick auf 10 Jahre Workshop „Karriere in der Allgemeinmedizin.“ *Z Allg Med.* 2018; 94(12): 500-505

Quinzler R, Freitag MH, **Wiese B**, Beyer M, Brenner H, Dahlhaus A, Doring A et al. A novel superior medication-based chronic disease score (medCDS) predicted all-cause mortality in independent geriatric cohorts. *J Clin Epidemiol.* 2019; 105: 112-124

Rihl M, Baerlecken N, **Wiese B**, Schmidt RE, Zeidler H. Intravenous Glucocorticoid Pulse Therapy in Active, NSAID Refractory Axial Ankylosing Spondylitis: A Retrospective Analysis Spanning 12 Months. *J Arthritis.* 2018; 7(1): 266

Schäfer I, Kaduszkiewicz H, Mellert C, Löffler C, Mortsiefer A, Ernst A, Stolzenbach C-O, **Wiese B** et al. Narrative medicine-based intervention in primary care to reduce polypharmacy: results from the cluster-randomised controlled trial MultiCare AGENDA. *BMJ Open.* 2018; 8(1): e017653

**Schneider N**, Egidi G, Jonitz G. Pharmaindustryunabhängige Aus-, Weiter- und Fortbildung. In: Lieb K, Klemperer D, Kölbel R et al. (Eds.) *Interessenkonflikte, Korruption und Compliance im Gesundheitswesen.* Berlin: Medizinisch Wissenschaftliche Verlagsgesellschaft; 2018. S.229-236

Stark A, Kaduszkiewicz H, Stein J, Maier W, Heser K, Weyerer S, Werle J, **Wiese B, Mamone S** et al. A qualitative study on older primary care patients' perspectives on depression and its treatments - potential barriers to and opportunities for managing depression. *BMC Fam Pract.* 2018; 19: 2

**Stiel S.** Allgemeiner Teil: Dokumentation und Assessment. In: Sabatowski R, Maier BO, Ostgathe C et al. (Eds.) Palliativmedizin – 1000 Fragen: Für Klinik, Praxis und die Zusatzweiterbildung Palliativmedizin. Stuttgart: Georg Thieme Verlag. 2., aktualisierte Auflage: 24-26

**Stiel S**, Heckel M, Wendt KN, Weber M, Ostgathe C. Palliative Care Patients' Quality of Dying and Circumstances of Death—Comparison of Informal Caregivers' and Health-Care Professionals' Estimates. *Am J Hosp Palliat Med.* 2018; DOI: 10.1177/1049909118756616

**Stiel S**, Nurnus M, Ostgathe C, Klein C. Palliative sedation in Germany: factors and treatment practices associated with different sedation rate estimates in palliative and hospice care services. *BMC Palliat Care.* 2018; 17(1): 48

**Stiel S**, Stelzer E-M, **Schneider N**, **Herbst FA**. Exploring end-of-life interaction in dyads of parents and adult children: a protocol for a mixed-methods study. *BMC Palliat Care.* 2018; 17(1): 68

Tiedtke JM, **Stiel S**, Heckel M, **Herbst FA**, Sturm A, Sieber C et al. Staff members' ambivalence on caring for patients with multidrug-resistant bacteria at their end of life: A qualitative study. *J Clin Nurs.* 2018; 27(15-16): 3115-3122

Werdecker L, Kujawa M, **Schneider N**, Neitzke G. Advance care planning: Eine qualitative Untersuchung zur Patientenperspektive bei fortgeschrittener Herzinsuffizienz. *Z Gerontol Geriatr.* 2018; DOI: 10.1007/s00391-018-1385-9